

661.566: 66.097.3:669.054.8

• • • • • , • • • • • , • • • • • , • • • • • ;
" " " " " , •

The possibility of the utilization of platinum containing wastes of nitric acid production is studied. Quantitative content of the wastes containing platinum group metals from the surface of boil-utilizator and absorption column is shown. In this article content of wastes with platinum group metals and choose sedimentator for metal platinum obtaining are made.

,
 -
 ,
 ,
 ,
 -
 -
 .
 0,716 ,
 - , , -
 , .
 ,
 ,
 " " ,
 " -
 , , -
 -
 . -
 .
 , -
 0,716 . 2400 -
 .
 -
 1.
 1

	, % .
	50,43
	1,37
	16,87
	1,1
	0,16
	0,34
	0,1
	17,75
	0,05
	0,25

	0,92
	0,16

,

,

-

2.

,

2

,

	, %
	11,35
	1,05
	9,98
	13,43
	4,21
	0,25
	0,03
	27,28
	0,18
	0,22
	0,06

. 1

. 2,

,

,

,

-

.

-

,

.

,

,

-

.

,

,

,

.

,

.

:

,

,

,

,

..

[5]

.

,

,

,

(

,

,

)

.


,

,

-

.

,



800...1000 °C

~ 10 %

[6].

[7].

: 1.

1985. – 327 . 2.

/

, 1999 – 368 . 3.

. – . 1983. – 286 . 4.

, 1974. - 17. - . 10. - . 1587 – 1589. 5.

, 1972 – 613 .

6.

//

2

"

". – . – 1997. – . 1. – . 189-190. 7.

.: , 1965 – 215 .

16.04.07